

REFERENCES

- Albee, A.L., 1962, Relationships between the mineral association, chemical composition and physical properties of the chlorite series: Am. Mineralogist, 47, 851-870.
- Ames, L.L. Jr., 1961, Volume relationships during replacement reactions: Econ. Geol., 56, 1438-1445.
- Ames, R.L., 1962, Sulphur isotope study of the Tintic mining districts, Utah: Unpublished Ph.D. dissertation, Yale University.
- Aw, P.C., 1967. Ignimbrites in central Kelantan, Malaya: Geol. Mag., 104, 13-18.
- Babcock, R.S., 1973, Computational models of metasomatic processes: Lithos, 6, 279-290.
- Bachinski, D.J., 1969, Bond strength and sulphur isotopic fractionation in coexisting sulphides: Econ. Geol., 64, 56-65.
- Barnes, H.L., and Czamanske, G.K., 1968, Solubilities and transport of ore minerals: In Geochemistry of Hydrothermal Ore Deposits, H.L. Barnes (ed.), Holt, Rinehart and Winston, Inc., New York, p.334-381.
- Bastin, E.S., 1950, Interpretation of ore textures: Geol. Soc. Amer. Mem., 45, 101p.
- Bath, T.F.W., 1948, Oxygen in rocks: A basis for petrographic calculations: Jour. Geology, 56, 50-60.
- _____, 1959, Principles of classification and norm calculations of metamorphic rocks: J. Geol., 67, 135-152.

- Becker, R.H., and Clayton, R.N., 1972, Carbon isotopic evidence for the origin of a banded iron-formation in Western Australia: Geochim. et Cosmochim. Acta, 36, 577-596.
- Berry, L.G., 1940, Studies of mineral sulphosalts IV - Galenobismutite and "Lillianite": Univ. Toronto, Geol. Ser., 44, 5-19.
- _____, and Thompson, R.M., 1962, X-ray powder data for ore minerals: the Peacock Atlas: Geol. Soc. Amer., Mem. 85.
- Bishop, D.G., 1972, Transposition structures associated with cleavage formation in the Otago schists: N.Z. Jour. Geol. Geophys., 15, 360-371.
- Blank, R.H., 1962, Ignimbrite vent in the Bull Valley District, Utah (abs.): Geol. Soc. Amer., spec. paper 73, 27p.
- Bodechtel, J. and Klemm, D.D., 1966, Über die lagerstättenkundliche Stellung und chemische Zusammensetzung der Bleiwismutspiessglanze. Geol. Rundschau., 55, 418-427.
- Borisenko, L.F., Lebedeva, S.I., and Serdobova, L.I., 1969, Titanium magnetite and magnetite in iron ore deposits of different genesis: Int. Geol. Rev., 11, 1408-1418.
- Boydell, H.C., 1925, The role of colloidal solutions in the formation of mineral deposits. Bull. Inst. Min. Metall. 243, 1-108.
- Brown, G., 1955, The effect of isomorphous substitutions on the intensities of 001 reflections of mica and chlorite type structures. Mineral Mag., 30, 657-665.
- Burst, J.F. Jr., 1969, Diagenesis of Gulf Coast Clayey sediments and its possible relation to petroleum migration: Am. Assoc. Petrol. Geo. Bull., 53, 73-93.

- Cherapanov, V.A., 1951, Aspects of morphology structure and substitution of malachite aggregates from Urals, deposits: Zap.Vses. Min. obs., 18, 3, 214-219. (In Russian, quoted by Chukhrov, 1965)
- Chukhrov, F.V., 1965, Present views on colloids in ore formation. Int. Geol. Rev., 8, 3, 336-346.
- _____, 1973, On the genesis problem of thermal sedimentary iron ore deposits: Mineralium Deposita, 138-147.
- Clayton, R.N., Friedman, I., Graf, D.O., Mayeda, T.A., Meets, W.F., and Shimp, N.F., 1966, The origin of saline formation waters I. Isotopic composition: J. Geophys. Res., 71, 3869-3882.
- Cook, H.E., 1968, Ignimbrite flows, plugs and dykes in the southern part of the Hot Creek Range, Nye County, Nevada: Geol. Soc. Amer. Mem., 116, 107-152.
- Craig, H., 1963, The isotopic geochemistry of water and carbon in geothermal areas: In Nuclear Geology in Geothermal Areas, E. Tongiorgi (ed.), Consiglio Nazionale delle Ricerche, Pisa, p.17-53.
- _____, 1966, Isotopic composition and origin of the Red Sea and Salton Sea geothermal brines: Science, 154, 1544-1548.
- Craig, J.R., 1967, Phase relations and mineral assemblages in the Ag-Bi-Pb-S system: Mineralium Deposita, 1, 278-306.
- Crohn, P.W., and Oldershaw, W., 1965, The Geology of the Tennant Creek One Mile Sheet Area, N.T.: Bur. Min. Resour. Aust. Rep. 83, 72p.
- Davidson, A., and Wyllie, P.J., 1965, Zoned magnetite and platy magnetite in Cornwall type ore deposits: Econ. Geol., 60, 766-771.

- Deer, W.A., Howie, R.A., and Zussman, J., 1966, An introduction to the rock-forming minerals, Longmans, London, 528p.
- Degens, E.T., and Epstein, S., 1962, Relationship between O^{18}/O^{16} ratios in coexisting carbonates, cherts, and diatomites: Amer. Assoc. Petrol. Geol. Bull., 46, 534-542.
- Dickey, P.A., 1966, Patterns of chemical composition in deep subsurface waters: Am. Assoc. Petrol. Geol. Bull., 57, 265-273.
- Drummond, A.D., Trotter, J., Thompson, R.M., and Gower, J.A., 1969, Neyite a new sulphosalt from Alice Arm. British Columbia. Can. Mineralogist, 10, 90-96.
- Duffell, S., 1937, Diffusion and its relation to ore deposition: Econ. Geol., 32, 494-510.
- Duncan, D.McP., 1970, Selected porphyroidal and granitic rocks at Tennant Creek, Northern Territory: Unpublished Ph.D. thesis, University of Tasmania.
- Dunnet, D., and Harding, R.R., 1967, Geology of the Mount Woodcock 1-mile Sheet Area, Tennant Creek, N.T.: Bur. Min. Resour. Aust. Rep., 114, 57p.
- Dunoyer de Segonzac, G., 1970, The transformation of clay minerals during diagenesis and low grade metamorphism: A review: Sedimentology, 16, 281-356.
- D'yachkova, I.B., and Khodakovskiy, I.L., 1968, Thermodynamic equilibria in the systems S-H₂O, Se-H₂O, and Te-H₂ in the 25-300°C temperature range and their geochemical interpretations: Geochem. Int., 5, 1108-1125.

- Dymkin, A.M., and Sokolov, G.A., 1961, Colloform segregations of endogenous magnetite in the Kurshunkul' Deposits. Geol. i geofiz., 1, 77-85 (In Russian).
- Edwards, A.B., 1955, The composition of the Peko Copper Orebody, Tennant Creek: Aust. Inst. Min. Met. Proc., 175, 55-82.
- Ellis, A.J., 1967, The chemistry of some explored geothermal systems: in Geochemistry of Hydrothermal Ore Deposits, H.L. Barnes (ed.), Holt, Rinehart and Winston, New York, 465-514.
- Elliston, J.N., 1960, Ore localisation by Preconsolidation Structures: Aust. Inst. Min. Met. Proc., 196, 29-50.
- _____, 1963, Sediments of the Warramunga Geosyncline. In Syntaphral Tectonics and Diagenesis: "Symposium". (S.W. Carey Convener), Geol. Dept. Univ. Tasmania, L1-L45.
- _____, 1966, The Genesis of the Peko Orebody: Aust. Inst. Min. Met. Proc., 218, 9-18.
- _____, 1968, Retextured Sediments: Intern. Geol. Congr. 23rd, Rept. Session, Pragne, 8, 85-104.
- _____, 1969, The genesis of some epigenetic type ore deposits: Cas. Mineral. Geol., 14, 129-139.
- Engel, A.E.J., Clayton, R.N., and Epstein, S., 1958, Variations in isotopic composition of oxygen and carbon in Leadville limestone and in its hydrothermal and metamorphic phases: Jour. Geol., 66, 374-393.
- Esquevin, R., 1969, Influence de la composition chimique des illites sur leur cristallinite: Bull. Centre. Rech. Pav.-S.N.P.A., 3, 147-154.

- Eugster, H.P., and Wones, D.R., 1962, Stability relations of the ferruginous biotite, annite: J. Petrology, 3, 82-125.
- Fander, H.W., 1966, Selenium at Peko and Cobar: AMDEL Bulletin (Australia), 2, 73-75
- Fawcett, J.J., and Yoder, H.S., 1966, Phase relationships of chlorites in the system $MgO-Al_2O_3-SiO_2-H_2O$: Am. Mineralogist, 51, 353-380.
- Fink, G., 1910, Bidrag till Sveriges Mineralogi: Ark. Kemi, Mineral, Geol., 3, No. 35, p.4.
- Fischer, A.G., 1964, The lofer cyclothems of the Alpine Triassic, Bull. Kansas Geol. Surv., 169, 108-149.
- Forbes, W.C., 1969, Unit-cell parameters and optical properties of talc on the join $Mg_3Si_4O_{10}(OH)_2 - Fe_3Si_4O_{10}(OH)_2$: Am. Mineralogist, 54, 1399-1408.
- _____, 1971. Iron content of talc in the system $Mg_3Si_4O_{10}(OH)_2 - Fe_3Si_4O_{10}(OH)_2$: Jour. Geology, 79, 63-74.
- Fron del, C., 1938, Stability of colloidal gold under hydrothermal conditions: Econ. Geol., 33, 1-20.
- Garrels, R.M., 1959, Rates of geochemical reactions at low temperatures and pressures: Researches in Geochemistry, V.I, P.A. Abelson (ed.), Wiley, New York.
- _____, and Christ, C.L., 1965, Solutions, Minerals and Equilibria, Harper and Row, New York, 450p.
- _____, Dreyer, R.M., and Howland, A.L., 1949, Diffusion of ions through intergranular spaces in water-saturated rocks: Geol. Soc. Amer. Bull., 60, 1809-1828.

- Goldschmidt, V.M., and Strock, L., 1935, Zur Geochemie der Selen. II: Nachr. Ges. Wiss. Gottingen, Math-Nat. Kl., N.F. Fachgr., 1, 123-142.
- Goldsmith, R.J., and Newton, R.C., 1969, P-T-X relations in the system $\text{CaCO}_3\text{-MgCO}_3$ at high temperatures and pressures: Am. J. Sci., 276A 160-190.
- Goleva, G.A., and Lushnikov, V.V., 1967, Occurrence of Selenium in the groundwaters of ore deposits and in some types of mineral waters: Geochem. Int., 4, 378-385.
- Goni, J., Guillemin, C., and Sarcia, C., 1967, Geochimie der l'or exogene - Etude experimentale de la formation des dispersions colloïdales d'or et de leur stabilite: Mineralium Deposita, 1, 259-268.
- Goodwin, A.M., 1956, Facies relations in the Gunflint iron formation: Econ. Geol., 51, 565-595.
- Goulevitch, J., (in press, April 1974), Geology and Structural Control at the Warrego Mine: in Economic Geology of Australia and Papua New Guinea (Metalliferous Volume): Melbourne, Aust. Inst. Min. Metal.
- Graham, A.R., 1950, Synthesis and X-ray study of compounds in the systems Pb-Bi-S, Ag-Bi-S and Ag (Bi,Sb₂)S₂. Unpublished Ph.D. Thesis, Univ. of Toronto. 100p.
- _____, Thompson, R.M., and Berry, L.G., 1953, Studies of mineral sulphosalts: XVII: Cannizzarite: Am. Mineralogist, 38, 536-544.
- Grigor'ev, D.P., 1965, Ontogeny of Minerals: Israel Program for Scientific Translations, Jerusalem, 250p.

- Grinenko, V.A., and Grinenko, L.N., 1967, Fractionation of sulphur isotopes in high temperature decomposition of sulphides by water vapour: Geochemistry Int., 4, 843-848.
- Grip, E., and Wirstam, A., 1970, The Boliden Sulphide deposit, A review of geo-investigations carried out during the lifetime of the Boliden Mine, Sweden (1924-1967). Sver. Geol. Unders., Ser. C, No. 651.
- Gross, G.A., 1970, Nature and occurrence of iron deposits. In Survey of World Iron Ore Resources, United Nations Dept. Econ. Social Affairs, New York.
- Gruner, J.W., 1934, The crystal structure of talc and pyrophyllite: Zeit. Krist., 88, 412-419.
- _____, 1944, The composition and structure of Minnesotaite, A common iron silicate in iron formations: Am. Mineralogist, 29, 363-372.
- Hall, W.E., and Friedman, I., 1963, Composition of fluid inclusions, Cave-in-Rock fluorite district, Illinois, and Upper Mississippi Valley, Zinc-lead district: Econ. Geol., 58, 886-911.
- _____, 1969, Oxygen and carbon isotopic composition of ore and host rock of selected Mississippi Valley deposits: U.S. Geol. Survey Prof. Paper, 650-C, C140-C148.
- Haranczyk, Czeslaw, 1969, Non colloidal origin of colloform textures, (discussion): Econ. Geol., 64, 466-468.
- Harris, D.C., and Owens, D.R., 1973, Berryite, a Canadian occurrence. Can. Mineralogist, 11, 1016-1018.

- Healey, T.W., 1972, Research in colloid and surface chemistry - Physiochemical processes in the diagenesis of sediments: Unpublished report, University of Melbourne, Department of Physical Chemistry, 103p.
- Heinrich, K.F.J., 1966, X-ray absorption uncertainty. In: The Electron Microprobe, New York, J. Wiley & Sons, 296-377.
- Helgeson, H.C., 1964, Complexing and hydrothermal ore deposition, International series of Monographs on earth Sciences, Pergamon Press, Oxford, 128p.
- _____, 1967, Solution chemistry and metamorphism: in Researches in Geochemistry, V.II, Abelson, P.H. (ed.), John Wiley & Sons, New York, 362-404.
- _____, 1968, Geologic and thermodynamic characteristics of the Salton Sea geothermal system: Am. J. Sci., 266, 129-166.
- _____, 1969, Thermodynamics of hydrothermal systems at elevated temperatures and pressures: Am. J. Sci., 267, 729-804.
- _____, 1970, A chemical and thermodynamic model of ore deposition in hydrothermal systems. In, Fiftieth Anniversary Symposia: Mineralogy and Petrology of the Upper Mantle, Sulphides. Mineralogy and Geochemistry of Non-Marine Evaporites. Mineral. Soc. Amer. Spec. Pap. 3, 155-186.
- _____, Brown, T.H., and Leeper, R.H., 1969, Handbook of Theoretical Activity Diagrams Depicting Chemical Equilibria in Geologic Systems Involving Minerals and an Aqueous Phase at one atm. and 0°-300°C., Freeman, Cooper, and Co., San Francisco, 253p.

- _____, and Garrels, R.M., 1968, Hydrothermal transport and deposition of gold: Econ. Geol., 63, 622-635.
- Hemley, J.J., and Jones, W.R., 1964, Chemical aspects of hydrothermal alteration with emphasis on hydrogen metasomatism: Econ. Geol., 59, 538-569.
- Henley, K.J., 1971, Mineralogical investigation of bismuth losses in Juno testing. Unpublished Amdel Report, (No. 750).
- _____, Radke, F., and Schultz, P.K., 1971, Mineralogical investigation of seven samples of Juno concentrates. Unpublished Amdel Report, (No. 820).
- _____, and Schultz, P.K., 1972, The Bi-Pb-(Cu)-S-Se Mineral in the Juno orebody, Tennant Creek. Paper presented at Specialist Meeting of Geol. Soc. Aust., February, 1972.
- Hey, M.H., 1954, A new review of the chlorites: Mineral Mag., 30, 277-292.
- Holser, W.T., 1947, Metasomatic processes: Econ. Geol., 42, 384-395.
- _____, and Schneer, C.J., 1961, Hydrothermal magnetite: Geol. Soc. Amer. Bull., 72, 369-386.
- Irving, J.D., 1911, Replacement ore-bodies and the criteria for their recognition: Econ. Geol., 6, 527-561, 619-669.
- Ivanac, J.F., 1954, The Geology and Mineral Deposits of the Tennant Creek Gold-Field, N.T.: Bur. Min. Resour. Aust. Bull. 22, 164p.
- Ivanov, S.N., 1969, Factors in the Genesis of Hydrothermal Ore Deposits: Dokl. Akad. Nauk. SSSR., 186, 30-32.

- James, H.L., 1954, Sedimentary facies of iron formation: Econ. Geol., 49, 235-293.
- _____, 1966, Chemistry of the iron-rich sedimentary rocks, U.S. Geol. Surv. Prof. Pap., 440-W, 61p.
- Jensen, M.L., 1965, The rational and geological aspects of solid diffusion, Can. Mineralogist, 8, 271-290.
- Jensen, M.L., 1967, Sulphur isotopes and mineral genesis: in, Geochemistry of Hydrothermal Ore Deposits; H.L. Barnes (ed.), Holt, Rinehart and Winston, New York, 143-165.
- Jirgensons and Straumanis, 1954, A short textbook on Colloid Chemistry, Pergamon Press, Oxford. 500p.
- Johansson, K., 1924, Bidrag till Gladhammar-gruvornas mineralogi. Ark. Kemi, Mineral., Geol., 9, No. 8, 1-22.
- _____, 1924a, Ett par selenförändringar mineral från Falu gruva. Ark. Kemi, Mineral., Geol., 9, No. 8, 22-28.
- Jones, G.J., 1970, The geology of Gecko Mine, Tennant Creek: paper presented at Geopko Technical Seminar, King Island, 1970.
- Jones, P.H., 1968, Hydrodynamics of geopressure in the northern Gulf of Mexico Basin: Amer. Inst. Min. Eng. Mtg. Spec. Pap., 2207, 12.
- Kalocsai, G.I.Z., (in prep.), Precipitation of gold from aqua regia solutions and subsequent determination by atomic absorption.
- Kaplan, I.R., Emery, K.O., and Rittenberg, S.C., 1963, The distribution and isotopic abundance of sulphur in recent marine sediments off southern California: Geochim. et Cosmochim. Acta, 27, 297-331.

- Karup-Møller, S., 1966, Berryite from Greenland, Can. Mineralogist, 8, 414-423.
- _____, 1970, Gustavite, a new sulphosalt mineral from Greenland, Can. Mineralogist, 19, 175-190.
- _____, 1970a, Weibullite, laitakarite and bismuthinite from Falun Sweden. Geol. Foeren, Stockholm, Foerh., 92, 181-187.
- _____, 1972, New data on pavonite, gustavite, and some related sulphosalt minerals: Neues Jahrb. Mineral., Abh., 117, No. 1, 19-38.
- Kelso, P., (in press), Bismuth backgrounds, Tennant Creek, Northern Territory: Jour. Geochem. (Aust.)
- Keith, M.L., and Weber, J.N., 1964, Carbon and oxygen isotopic composition of selected limestones and fossils: Geochim. et Cosmochim. Acta, 28, 1787-1816.
- Kennedy, G.C., 1950, A portion of the system silica-water: Econ. Geol., 45, 629-653.
- Kharaka, Y.K., Berry, F.A.F., and Friedman, I., 1973, Isotopic composition of oil-field brines from Kettleman North Dome, California, and their geologic implications: Geochim. et Cosmochim. Acta, 37, 1899-1908.
- Kingston, P.W., 1968, Studies of mineral sulphosalts: XXI - Nuffieldite, a new species: Can. Mineralogist, 9, 439-452.
- Kitahara, S., 1960, The solubility of quartz in water at high temperatures and high pressures: Rev. Phys. Chem. Japan, 30, 109-114.

- Klominsky, J., Reider, M., Keift, C., and Mraz, L., 1971, Heyrovskyite, $6(\text{Pb}_{0.86}\text{Bi}_{0.08}(\text{Ag,Cu})_{0.04}) \text{S} \cdot \text{Bi}_2\text{S}_3$ from Hurky, Czechoslovakia, a New Mineral of Genetic interest. Mineralium, Deposita, 6, 133-147.
- Klyakhin, V.A., and Dmitriyeva, M.T., 1968, More information about synthetic and natural lillianite. Dokl. Akad. Nauk. SSSR, 178, 173-175.
- Kodera, M., Kupcik, V., and Makovicky, E., 1970, Hoderushite - a new sulphosalt: Mineral. Mag., 37, 641-648.
- Kolonin, G.R., and Aksenova, T.P., 1970, Investigation of complexing of some heavy metals in NaCl solutions at 20-90°C: Geochem. Int., 11, 973-978.
- Kopeykin, V.A., and Mikhaylov, A.S., 1970, Solubility and forms of occurrence of silica in normal dilute solutions: Dokl. Akad. Nauk, SSSR, 191, 193-195.
- Korzhinskii, D.S., 1955, An outline of metasomatic processes: in Basic problems in the study of magmatogenic deposits, Academy of Sciences of the U.S.S.R. (Translated in Int. Geol. Rev., 6, 1964; 1713-1734, 1920-1952 and 2169-2198.)
- _____, 1957, Acidity conditions in post-magmatic solutions: Izv. Akad. Nauk. SSSR., Ser. Geol., 12, 3-12 (in Russian).
- _____, 1959 The advancing wave of acidic components in ascending solutions and hydrothermal acid-base differentiation: Geochim. et Cosmochim. Acta, 17, 17-20.
- _____, 1964, A theory of the process of mineral formation: Int. Geol. Rev., 6, 387-398.

- _____, 1968, The theory of Metasomatic Zoning: Mineralium Deposita, 3, 222-231.
- _____, 1970, Theory of Metasomatic Zoning, Clarendon Press, Oxford, 162p.
- Kostov, I., 1958, Bonchevite, PbBi_4S_7 , a new mineral: Mineral. Mag., 31, 821-828.
- Krauskopf, K.B., 1951, The solubility of gold: Econ. Geol., 46, 858-870.
- _____, 1956, Dissolution and precipitation of silica at low temperatures: Geochim. et Cosmochim. Acta, 10, 1-26.
- _____, 1964, Possible role of volatile metal compounds in ore genesis. Econ. Geol., 59, 22-45.
- Lakin, H.W., 1965, Geochemistry of selenium in relation to agriculture: U.S. Dept. Agr. Handbk., 200, 3-12.
- Landa, E.R., and Gast, R.G., 1973, Evaluation of crystallinity in hydrated ferric oxides: Clays and Clay Minerals, 21, 121-130.
- Lapukhov, A.S., 1966, Diffusion transfer of material in fissures and through the intergranular spaces of water-saturated rocks: Geologiya Geofiz. Novosibirsk, 12, 56-64 (in Russian).
- Large, R.R., 1971, Metasomatism and scheelite mineralisation at Bold Head, King Island: Aust. Inst. Min. Met. Proc., 238, 31-45.
- _____, 1971a, Regional Mapping in the Queen of Sheba, Three Ways, Whippet Area: Unpublished company Report, Geapeko, Ltd.

- _____, 1972, Metal Zonation at Tennant Creek and its use as a guide in surface and underground exploration: Unpublished Company Report, Geopeko Ltd.
- Lasky, S.G., 1930, A colloidal origin of some of the Kennecott ore minerals: Econ. Geol., 25, 737-757.
- Lebedev, L.M., 1967, Metacolloids in endogenic deposits. Monographs in Geoscience, R.W. Fairbridge (ed.), Plenum Press, New York, 298p.
- Levorsen, A.L., 1967, Geology of Petroleum: W.H. Freeman and Co., San Francisco, 723p.
- Lindgren, W., 1912, The nature of replacement: Econ. Geol., 7, 521-535.
- _____, 1925, Gel replacement a new aspect of metasomatism. Nat. Acad. Sci., Proc., 11, 5-11.
- _____, 1933, Mineral Deposits, McGraw-Hill, New York, 930p.
- Lindstrom, G., 1887, Title Unknown: Geol. Foeren, Stockholm Foerh., 9, 523.
- Lovering, T.S., 1949, Rock alteration as a guide to ore - East Tintic District, Utah: Econ. Geol., Monograph 1, 64p.
- Mackenzie, R.C., and Meldan, K., 1959, The ageing of sesquioxide gels.
(1) Iron Oxide gels: Mineral Mag., 32, 153-165.
- Makovicky, E., and Maclean, W.H., 1972, Electron Microprobe analysis of Hodrushite: Can. Mineralogist, 11, 504-513.
- McCrea, J.M., 1950, The isotopic chemistry of carbonates and a paleo-temperature scale: Jour. Chem. Physics, 18, 849-857.

- McNeil, R.D., 1963, Slip complexes in the Warramunga Geosyncline: in Syntaphral Tectonics and Diagenesis. "Sumposium". (S.W. Carey Convener), Geol. Dept. Univ. Tasmania, M1-M11.
- _____, 1966, Geology of the Orlando Mine, Tennant Creek, Australia: Econ. Geol., 61, 221-242.
- Malashin, V.A., 1968, Differentiation of matter in the stage of dispersive metasomatism: Dokl. Akad. Nauk. SSSR., 182, 1406-1409.
- Marakushev, A.A., and Perchuk, L.L., 1970, On the influence of the acidity and temperature of postmagmatic solutions on the composition of micas and chlorite, Union. Geol. Sci. A. No. 2: Problems of Hydrothermal Ore Deposition, Schweizerbart, Stuttgart, 274-278.
- Marowsky, G., and Wedepohl, K.H., 1971, General trends in the behaviour of Cd, Hg, Tl and Bi in some major rock forming processes: Geochem. et Cosmochim. Acta, 35, 1255-1267.
- Martin, R.F., and Piwinski, A.J., 1969, Experimental data bearing on the movement of iron in an aqueous vapor: Econ. Geol., 64, 798-803.
- Maxwell, J.C., 1962, Origin of slaty and fracture cleavage in the Delaware Water Gap area, New Jersey and Pennsylvania. In Engel, A.E.J., Jones, H.L., and Leonard, B.F., (eds.), Petrologic Studies: A volume in honour of A.F. Buddington: Geol. Soc. Amer., 660p.
- Meyer, C., and Hemley, J.J., 1968, Wall Rock Alteration: in, Geochemistry of hydrothermal ore deposits, H.L. Barnes (ed.), Holt, Rinehart and Winston, New York, 166-235.
- Moench, R.H., 1966, Relation of S_2 schistosity to metamorphosed clastic dykes, Rangeley - Phillips area, Maine: Geol. Soc. Amer. Bull., 77, 1449-1462.

- Milsom, J.S., and Finney, W.A., 1965, Tennant Creek Detailed Aeromagnetic Survey, Northern Territory: Bur. Min. Resour. Aust. Rec. 1965/50 (unpubl.).
- Minster, E.F., 1967, Benjaminite - $(\text{Cu}, \text{Ag})_2\text{Pb}_2\text{Bi}_4\text{S}_9$: Dokl. Akad. Nauk. SSSR, 194, 675-678.
- Moore, E.S., 1932, A magnetite vein of abnormal type: Econ. Geol., 27, 387-390.
- Moore, P.B., 1967, A classification of sulphosalt structures derived from the structure aikinite: Am. Mineralogist, 52, 1874-1876.
- Morse, H.W., Warren, C.H., and Donnay, J.D.H., 1932, Artificial spherulites and related aggregates: Am. J. Sci., 23, 421-439.
- Naboko, S.I., 1959, Volcanic exhalations and products of their reactions as exemplified by the Kamchatka-Kuriles volcanoes: Bull. Volcanol., 20, 121-136.
- Nockolds, S.R., 1966, The behaviour of some elements during fractional crystallisation of magma: Geochim et Cosmochim. Acta., 30, 267-278.
- Northrop, D.A., and Clayton, R.N., 1966, Oxygen-isotope fractionations in systems containing dolomite: Jour. Geol., 74, 174-196.
- Nowacki, W., and Stalder, H.A., 1969, Zwei Wismutsulfosalze von Sta. Maria. Val. Medel, Mt. Graubunden, Schweiz, Schweiz Mineral. Petrog. Mitt., 49, 97-101.
- Nuffield, E.W., and Harris, D.C., 1966, Studies of mineral sulphosalts: XX: Berryite, a new species: Can. Mineralogist, 8, 407-413.

- Odman, O.H., 1941, Geology and Ores of the Boliden deposit, Sweden.
Sver., Geol. Unders., Ser. C, No. 438,
- Oertel, A.C., 1970. The Calculation of results in electron-probe micro-analysis. C.S.I.R.O. Aust. Div. Soils Tech. Pap., 9, 20p.
- Ohmura, M., and Nowacki, W., 1970, Note on the space group and on the structure of aikinite derivatives: Neues Jahrb. Mineral. Monatsh., 1970, 158-162.
- Ohmoto, Hiroshi, 1972, Systematics of sulphur and carbon isotopes in hydrothermal ore deposits: Econ. Geol., 67, 551-578.
- Ontoev, D.O., Troneva, N.V., and Tsepin, A.I., 1972, Argentiferous wittite from eastern Trans-Baikal: Zap. Vses. Mineral. Obshchest., 101. 476-480 (in Russian).
- Ontoyev, D.O., 1959, Colloform magnetite and the distinctive platy and tabular pyrite segregations in magnetite ores: Tr. Mineral, muzei. Akad. Nauk, SSSR., 9, 74-85 (In Russian).
- O'Neil, J.R., Clayton, R.N., and Mayeda, T., 1969, Oxygen isotope fractionation in divalent metal carbonates: Jour. Chem. Physics, 51, 5547-5558.
- Otto, H.H. and Strunz, H., 1968, Zur Kristallchemie Synthetischer Blei-Wismut-Spiessglanze. Neues Jahrb. Mineral., Abh., 108, 1-19.
- Padera, K., 1956, Beitrag zur Revision der Mineralien aus der Gruppe von Wismutglanz und aikinit: Chem. Erde, 18, 14-18.

- Papson-McGugan, J.E., 1970, The diagenesis and depositional environment of the Permian Ranger Canyon and Mowitch Formations, Ishbel Group, from the Southern Canadian Rocky Mountains: Sedimentology, 15, 363-418.
- Park, C.F., and MacDiarmid, R.A., 1970, Ore Deposits, 2nd Edition, W.H. Freeman, San Francisco.
- Pavlov, N.V., 1958, Problems of the genesis of the Endogenetic Magnetite ores of the Fungusska Syncline on the Siberian Platform: Izv. USSR. Acad. Sci., Ser. Geol., (1958), 9, 1-17.
- Peacock, M.A., and Berry, L.G., 1940, Rontgenographic Observations on ore minerals. Univ. Toronto, Geol. Ser., 44, 47-69.
- Perry, D.V., 1969, Skarn genesis at the Christmas Mine, Gila County, Arizona: Econ. Geol., 64, 255-271.
- Perry, E.A., and Hower, J., 1972, Late stage dehydration in deeply buried pelitic sediments: Am. Assoc. Petrol. Geol. Bull., 56, 2013-2020.
- Pettijohn, F.J., 1963, Chemical Composition of Sandstones - excluding Carbonate and Volcanic Sands. U.S. Geol. Surv. Prof. Pap., 440-S, 21pp.
- Petruk, W., 1964, Determination of the heavy atom content in chlorite by means of the X-ray diffractometer: Am. Mineralogist, 49, 61-71.
- Pinckney, D.M., and Rye, R.O., 1972, Variation of O^{18}/O^{16} , C^{13}/C^{12} , Texture and mineralogy in altered limestone in the Hill Mine, Cave-in-District, Illinois: Econ. Geol., 67, 1-18.

- Plotnikov, V.I., 1959, Separation of Se^{IV} by means of iron hydroxide. Zavodskaya laboratoriya, 6, (in Russian).
- Poldervaart, A., 1953, Petrological calculations in metasomatic processes: Am. J. Sci., 251, 481-504.
- Pospelov, G.I., 1966, The types and mechanisms of metasomatism: Int. Geol. Rev., 9, 890-898.
- Powell, C.McA., 1969, Intrusive sandstone dykes in the Siamo Slate near Negamee, Michigan: Geol. Soc. Amer. Bull., 80, 2585-2594.
- _____, 1972, Tectonically dewatered slates in the Ludlovian of the Lake District, England: Geol. Jour., 8, 95-110.
- Ramdohr, Paul, 1969, The ore minerals and their intergrowths. English trans., (Amstutz C.H.R., Ed.) Pergannon Press, 1174p.
- Randal, M.A., Brown, M.C., and Douth, H.F., 1966, Geology of the Helen Springs and Beetaloo 1:250,000 sheet areas, N.T., Bur. Min. Resour. Aust., 1966/110 (unpubl.)
- Ridge, J.D., 1968, Changes and Developments in Concepts of Ore Genesis, in Ore Deposits of the United States, 1933-1967, J.D. Ridge (ed.), 1713-1834.
- Roedder, E., 1967, Fluid inclusions as samples of ore fluids: in, Geochemistry of Hydrothermal Ore Deposits, H.L. Barnes (ed.), Holt, Rinehart and Winston, New York, 515-574.
- _____, 1968, The non colloidal origin of "colloform" textures in sphalerite ores: Econ. Geol., 63, 451-471.
- Rogers, A.F., 1917, A review of the amorphous minerals: Jour. Geology, 25, 515-541.

- Romberger, S.B., 1968, Solubility of copper in aqueous sulphide solutions coexisting with covellite from 25° to 200°C, with geologic applications: Unpublished Ph.D. Thesis, The Pennsylvania State Univ.
- Rundkvist, D.V., 1958, Morphological and internal structural features of mineral aggregates in near-surface tin ore deposits: Zap. Vses. Min. obs., 87, 423-439. (In Russian, quoted in Chukhrov, 1965.)
- Rust, G.E., 1935, Colloidal primary copper ores at Cornwall Mines, S.E. Missouri: Jour. Geology, 43, 398-426.
- Rye, R.O., 1966, The carbon, hydrogen, and oxygen isotope composition of the hydrothermal fluids responsible for the lead-zinc deposits at Providencia, Zacatecas, Mexico: Econ. Geol., 61, 1399-1427.
- Sakai, H., 1968, Isotopic properties of sulphur compounds in hydrothermal processes, Geochim. J. (Nagoya), 2, 29-49.
- Salanci, B., and Moh, G.H., 1969, Die experimentelle Untersuchung des pseudobinaren Schnittes $PbS-Bi_2S_3$ innerhalb des Pb-Bi-S-systems in Beziehung zu natürlichen Blei-Wismut-Sulfosalzen. Neues. Jahrb. Min., Abh., 112, 63-95.
- Sales, R.H. and Meyer, C., 1950, Interpretation of wall-rock alteration at Butte, Montana. Colo. Sch. Mines Q., 45, (1B) 261-273.
- Sangster, D.F., 1965, Colloform magnetite in a contact metasomatic iron deposit (discussion), Econ. Geol., 60, p.824-826.
- _____, 1972, Precambrian volcanogenic massive sulphide deposits in Canada: A review: Geol. Surv. Can. Pap. 72-22, 44pp.
- Schneiderholm, H., 1952, Erzmikroskopisches Praktikum: Stuttgart, 274p.

- Schoen, R., 1962, Semi-quantitative analysis of chlorite by X-ray diffraction: Am. Mineralogist., 47, 1384-1393.
- Seward, T.M., 1973, The complexes of gold and the transport of gold in hydrothermal ore solutions: Geochim. et Cosmochim. Acta., 37, 379-400.
- Shannon, E.V., 1964, Benjaminite, a new sulphosalt mineral of the Klaprotholite group: U.S. Nat. Mus. Proc., 65, No. 24, 1-9.
- Sheppard, S.M.F., Nielsen, R.L., and Taylor, H.P. Jr., 1971, Hydrogen and Oxygen isotope ratios in minerals from porphyry copper deposits: Econ. Geol., 66, 515-542.
- _____, and Schwarcz, H.P., 1970, Fractionation between carbon and oxygen isotopes and magnesium between coexisting metamorphic calcite and dolomite: Contr. Mineral. Petrol., 26, 161-198.
- Siever, R., 1962, Silica solubility, 0°-200°C, and the diagenesis of siliceous sediments: Jour. Geology, 70, 127-150.
- Sillen, L.G., 1964, Stability constants of metal-ion complexes. Sec. I. Inorganic ligands: Chem. Soc. London Spec. Pub., 17.
- Sindeeva, N.D., 1962, Mineralogy and types of Deposits of Selenium and Tellurium, Interscience Publishers, New York, 363p.
- Skinner, B.J., White, D.E., Rose, H.J., and Mays, R.E., 1967, Sulphides associated with the Salton Sea geothermal brine: Econ. Geol., 62, 316-330.
- Smith, F.G., and Kidd, D.J., 1949, Hematite-geothite relations in neutral and alkaline solutions under pressure; Am. Mineralogist, 34, 403-412.

- Springer, G., and Demirosoy, S., 1969. Bertrag Zur Clarung der Existenz von Klaprothit. Neues, Jahrb. Mineral., Monatsh., 32-37.
- Springer, G., 1971, The synthetic solid-solution series $\text{Bi}_2\text{S}_3\text{-BiCuPbS}_3$ (bismuthinite-aikinite): Neues Jahrb. Mineral., Monatsh., 1971, 19-24.
- Stanton, R.L., and Rafter, T.A., 1966, The isotopic constitution of sulphur in some stratiform lead-zinc sulphide ores: Mineralium Deposita, 1, 16-29.
- _____, 1972, Ore Petrology, McGraw-Hill Book Company, New York, 712p.
- _____, 1972a, A preliminary account of chemical relationships between sulphide lode and "banded iron formation" at Broken Hill, New South Wales: Econ. Geol., 67, 1128-1145.
- Stemprok, M., 1970, Continuity or Discontinuity of Processes leading to Wall-Rock Alteration in Problems of Hydrothermal Ore Deposition, Schweizrebart, Stuttgart 1970. (union Geol. Sci. A. No. 2.)
- Stevenson, J.S., and Jeffrey, W.G., 1964, Colloform magnetite in a contact metasomatic iron deposit: Econ. Geol., 59, 1298-1305.
- Strickland, C.S., (in press, April 1974), The Gecko Copper Mine, Tennant Creek: in Economic Geology of Australia and Papua New Guinea (Metalliferous volume): Melbourne, Aust. Inst. Min. Metal.
- Sweatman, T.R., and Long, J.V.P., 1969, Quantitative electron-probe microanalysis of rock-forming minerals: J. Petrol., 10, 332-379.
- Syritso, L.F., and Senderova, V.M., 1964, The problem of the existence of lillianite (in Russian): Zap. Vses. Mineral. Obshchest., 93, 468-471. (Ref. in: Mineral. Abstr. 17, 174-175).

- Takagi, J., and Takeuchi, Y., 1972, Structural principles of the crystal-line phases in the $\text{PbS-Bi}_2\text{S}_3$ system (Abstr), Acta Crystallographica, 28B, 566.
- Taylor, H.P. Jr., 1967, Oxygen isotope studies of hydrothermal mineral deposits: in, Geochemistry of Hydrothermal Ore Deposits, H.L. Barnes (Ed.), Holt, Rinehart and Winston, New York, 109-142.
- _____, Frechen, J., and Degens, E.T., 1967, Oxygen and carbon isotope studies of carbonatites from the Laacher See district, West Germany, and the Alno district, Sweden: Geochim. et Cosmochim. Acta, 31, 407-430.
- Tilling, R.I., Gottfried, D., and Rowe, J.J., 1973, Gold abundance in igneous rocks: bearing on gold mineralisation: Econ. Geol., 68, 168-186.
- Turner, F.J., 1968, Metamorphic Petrology, McGraw-Hill, New York, 403p.
- Toulmin, P., and Clark, S.P. Jr., Thermal aspects of ore formation: in, Geochemistry of hydrothermal ore deposits, H.L. Barnes (Ed.), Holt, Rinehart and Winston, New York, 437-464.
- Uytenbogaardt, W., and Burke, E.A.J., 1971, Tables for microscopic identification of ore minerals: Elsevier Publishing Company, Amsterdam, 430p.
- Van Everdingen, R.O., 1970, The Paint Pots, Kootenay National Park, British Columbia - acid spring water with extreme heavy-metal content: Can. Jour. Earth Sci., 7, 831-852.
- Van Hook, H.J., 1960, The ternary system $\text{Ag-Zn-Bi}_2\text{S}_3\text{-PbS}$: Econ. Geol., 55, 759-788.


- Walters, L.J. Jr., Claypool, G.E., and Choquette, P.W., 1972, Reaction rates and δO^{18} variation for the carbonate-phosphoric acid preparation method: Geochim. et Cosmochim. Acta, 36, 129-140.
- Weaver, C.E., and Beck, K.C., 1971, Clay water diagenesis during burial: how mud becomes gneiss: Geol. Soc. Amer. Spec. Pap., 134.
- Wehrenberg, J.P., and Silverman, A., 1965, Studies of base metal diffusion in experimental and natural systems: Econ. Geol., 60, 317-350.
- Weibull, M., 1885, Om selenhaltig galenobismutite from Falun grufva. Geol. Foeren, Stockholm Foerh., 96, 657-666.
- Weissberg, B.C., 1970, Solubility of gold in hydrothermal alkaline sulphide solutions: Econ. Geol., 65, 551-556.
- Welin, E., 1966, Notes on the mineralogy of Sweden. 5. Bismuth-bearing sulphosalts from Gladhammer, a revision: Ark. Mineral. Geol. 4, 377-386.
- Wells, N., Selenium content of soil-forming rocks: N.Z. Jour. Geol. Geophys., 10, 198-208.
- White, D.E., 1968, Environments of generation of some base-metal ore deposits: Econ. Geol., 63, 301-335.
- _____, and Waring, G.A., 1961, A review of the chemical composition of gases from volcanic fumaroles and igneous rocks. U.S. Geol. Surv. Prof. Pap., 424-C, C311-312.
- Whittle, A.W.G., 1966, The paragenesis and origin of the Tennant Creek Mineral Deposits. Unpub. Ph.D. thesis, University of Adelaide, 227p.

- Williams, B.T., 1970, The Gecko mineral channel: paper presented at Geopeko Technical Seminar, King Island, 1970.
- Willis, J.L., and Daly, M.R., 1972, Investigations within Jubilee No. 2 reserve 306, Tennant Creek, N.T. Geol. Surv. Rec., Mines Branch, Northern Territory Administration, Department of the Interior. 17pp.
- Wolfe, C.W., 1938, Cannizzarite and bismuthinite from Vulcano. Am. Mineralogist, 23, 790-798.
- Wright, K., 1969a, Textures from some epigenetic mineral deposits of Tennant Creek - Central Australia. Reprinted from meeting on remobilisation of ores and minerals, Cagliari, 1969, 219-251.
- _____, 1969b, The presence of uraninite in the mineral bodies of Tennant Creek field: Unpublished Company Report, Geopeko Ltd.
- _____, 1970, Geology and ore reserves of Ivanhoe Mine, Unpublished Company report, Geopeko Ltd.
- Zies, E.G. 1929, The Valley of Ten Thousand Smokes - I. The fumarolic incrustations and their bearing on ore deposition. Tech. Pap. Natl. Geogr. Soc., 1, No. 4, 62-79.
- Zelenov, K.K., 1960, Migration and accumulation of iron and aluminium in the volcanic areas of the Pacific Ocean. Izv. USSR. Acad. Sci. Ser. Geol., 8, 58-74.
- Zharikov, V.A., 1965, The possible significance of electrokinetic phenomena in geochemistry: in Problems of geochemistry, Khitarov (Ed.), Science Press.

APPENDIX APROCEDURE USED IN THE CALCULATION OF NORMATIVE MINERALOGY

Due to the varied mineralogical composition of the lode rocks and country rocks, and the widespread variation in the composition of the chlorites present in these rocks, the usual normative calculation methods were inappropriate, and the following procedures were adopted in their place.

(a) Sedimentary Rocks: applies to norms presented in Tables 3.2, 3.3, 4.5 and 9.2. All the normative minerals except chlorite were taken to have their 'ideal composition' (used in the calculation of epinorms, Bath, 1959). Chlorite was given the formula



In hematite bearing sediments free of magnetite, all the FeO, MnO and MgO was assumed to be present in chlorite. This allowed a ready calculation of the Fe+Mn/Fe+Mn+Mg ratio of the chlorite and determination of normative chlorite based on the composition given above. In magnetite bearing sediments free of hematite, FeO is present in both magnetite and chlorite. In such samples, the Fe₂O₃ content of the rock was used to calculate normative magnetite and the residual FeO was assumed to be in chlorite. Norms for rocks containing hematite, magnetite and chlorite could not be calculated unless a value for the Fe+Mn/Fe+Mn+Mg ratio of the chlorites was assumed (as is the case in col. 5, Table 3.3).

Normative albite was calculated directly from the Na₂O while normative apatite was calculated from the P₂O₅ content. Excess CaO not present in apatite or calcite was used to give normative anorthite, while normative calcite was calculated from the CO₂ content. K₂O and residual Al₂O₃ are present in both muscovite and K-feldspar whose normative values were calculated by solving the simultaneous equations:

$$\begin{aligned}\text{normative muscovite} &= \frac{(16.9 \times \text{wt.\% Al}_2\text{O}_3) - (8.4 \times \text{wt.\% K}_2\text{O})}{433} \\ \text{normative K-feldspar} &= \frac{\text{wt.\% Al}_2\text{O}_3 - 38.44 \text{ (norm. musc.)}}{18.35}\end{aligned}$$

The norms obtained by this method gave good totals ($100 \pm 1\%$) and agree with visual estimates recorded during the microscopic study of the sediments.

(b) Lode Rocks: applies to norms presented in Tables 4.1 and 9.3.

The chloritised sediments at Juno and Gecko are free of magnetite and their normative mineralogy was calculated by the method described above for the magnetite free sediments.

The mineralised sediments at Juno contain magnetite, hematite and chlorite but no quartz. This residual SiO_2 , Al_2O_3 and MgO left after the calculation of normative muscovite, albite and anorthite was considered to be present in chlorite. Assuming the formula given above, this allowed the calculation of normative chlorite and the percentage of FeO in chlorite. The remaining FeO and Fe_2O_3 were divided between magnetite and hematite.

The magnetite-chlorite lode rocks contain negligible muscovite and feldspars and thus normative chlorite was calculated directly from the total Al_2O_3 content of the rocks.

In the talc-magnetite zone, Al_2O_3 was assumed to be present only in chlorite, and the residual MgO was used to calculate normative talc. The latter mineral was given the formula $\text{Mg}_{5.5}\text{Fe}_{0.5}\text{Si}_8\text{O}_{20}(\text{OH})_4$ although microprobe data (section 4.3.2) indicates a variable iron content throughout the zone - a factor which probably contributed to the poor totals given by norms in Table 4.1. Normative pyrite which only occurs in measurable quantities in the talc zone was calculated from the sulphur analysis after allowing for the other metal sulphides. Normative magnetite

was calculated from residual FeO and normative hematite from residual Fe_2O_3 (after accounting for pyrite and magnetite).

The normative mineralogy of the dolomite zone was calculated in a similar manner to the other zones. Dolomite was split into four components (CaCO_3 , MgCO_3 , MnCO_3 and FeCO_3) and CO_2 was partitioned amongst these according to the residual amounts of the metal oxides.

APPENDIX BADDITIONAL GEOCHEMICAL DATA ON CARRAMAN FORMATION GREYWACKE AND SHALES

Analyses by Australian Mineral Development Laboratories

| JUNO | Cu | S | Se ppm |
|----------------------------|-----|------|--------|
| <u>UNALTERED SEDIMENTS</u> | | | |
| DDH 9/71/30-34 | 45 | 300 | <2 |
| 51-55 | 70 | 500 | <2 |
| 67-71 | 200 | 500 | <2 |
| 113-117 | 25 | 1600 | <2 |
| 276-280 | 5 | 450 | <2 |
| 381-385 | 5 | 450 | <2 |
| 451-455 | 15 | 650 | <2 |
| <u>LEACHED SEDIMENTS</u> | | | |
| DDH 9/71/137-141 | 5 | 300 | <2 |
| 164-168 | 35 | 1250 | <2 |
| 214-218 | <5 | 300 | <2 |
| EXPLORER 46 | | | |
| <u>UNALTERED SEDIMENTS</u> | | | |
| DDH 46/4/3/291 | 5 | 250 | <2 |
| 298 | 5 | 250 | <2 |
| 307 | 15 | 200 | <2 |
| 318 | 5 | 250 | <2 |
| 334 | 100 | 300 | <2 |
| 342 | 50 | 400 | <2 |
| 353 | 5 | 300 | <2 |
| 378 | 5 | 250 | <2 |
| 391 | 50 | 350 | <2 |
| 403 | 5 | 300 | <2 |
| WEST PEKO | | | |
| <u>UNALTERED SEDIMENTS</u> | | | |
| DDH 2A/1104 | 10 | 300 | <2 |
| 1496 | 70 | 250 | <2 |
| 1870 | 520 | 950 | <2 |
| 2099 | 10 | 300 | <2 |
| 2118 | 85 | 300 | <2 |
| 2204 | 110 | 300 | <2 |

APPENDIX C

ROCK SAMPLES LODGED AT UNIVERSITY OF NEW ENGLAND, AND GEOPEKO LTD., TENNANT CREEK.

| University N.E. Rock Number | Geopeko Ltd. | | Locality | Lithology |
|-----------------------------------|------------------------|----------------------------|-------------------------------|-----------------------------------|
| | Thin Section Number | Polished Section Number | | |
| R27787a | | GP 213 | <u>JUNO MINE</u> 700 level | Bismuth sulphosalts |
| R27788a | | | 700 level | Junite in magnetite-chlorite rock |
| R27789a | | | 700 level | Junite |
| R27790 | | | 700 level | Junite and "wittite" |
| R27791a | | GP 127 | 7/167/1** | Junite |
| R27792a | | GP 204 | 8/166/73 | Junite |
| R27793a | | GP 88 | 700 level | Aikinite member |
| R27794a | | GP 112 | 7/89/13 | Wittite |
| R27795a | | | 700 level | Wittite |
| R27796a | | GP 131 | 7/38/70 | Wittite |
| R27797a | | GP 173 | 7/26/125 | Heyrovskyite |
| R27798a | | | 7/92/33 | Aikinite members |
| R27799a | | | 7/25/130 | Aikinite + junite |
| R27800a | | GP 178 | 8/170/42 | Aikinite members |
| R27801a | | | 7/26/91 | Junite |
| R27802a | | | 700 level | Emplectite |
| | | | | Junite + aikinite |
| | | | <u>WHIPPET FORMATION</u> | |
| | GT 377 | | N.E. of Three Ways | Massive Sandstone |
| | GT 1307 | | Whippet Trig | |
| | | | Station | Massive Sandstone |
| | | GP 478 | Whippet Trig | |
| | | | Station | Massive Sandstone |

* Indicates a thin section in addition to hand specimen lodged at U.N.E.

a Indicates a polished section in addition to hand specimen lodged at U.N.E.

** 7/167/1 = 700 level, DDH No. 167, 1 foot from collar.

University N.E.

Geopeko Ltd.

Rock
NumberThin Section Polished Section
Number Number

Locality

Lithology

BERNBOROUGH FORMATION

| | | |
|--------|-----------------------|---------------------|
| GT 376 | Bernborough Area | Crystal-vitric tuff |
| GT 471 | North of Whippet Trig | Crystal-vitric tuff |
| GT 481 | Bernborough Area | Crystal-vitric tuff |
| GT 442 | North of Whippet Trig | Laminated rhyolite |
| GT 311 | Bishops Creek | Pumice rich tuff |

CARRAMAN FORMATION

| | | |
|---------|-----------------------------------|--------------------------------|
| R27803 | Explorer 86 | Magnetic siltstone |
| R27804 | Explorer 86 | Magnetic siltstone |
| R27805 | Explorer 46 | Greywacke |
| R27806 | Explorer 46 | Chloritic siltstone |
| R27807 | Explorer 46 | Greywacke/hematite shale |
| R27808 | West Peko | Hematite siltstone |
| R27809 | 3/4 km north of Golden Forty Mine | Peko porphyry |
| R27810* | as above | Peko porphyry |
| R27811* | as above | Peko porphyry |
| R27812* | as above | Peko porphyry |
| R27813* | as above | Peko porphyry |
| R27814* | as above | Peko porphyry |
| R27815* | as above | Peko porphyry |
| R27816* | as above | Siltstone overlying porphyry |
| R27817* | as above | Greywacke overlying porphyry |
| R27818* | Jubilee area | Baveno Porphyry |
| R27819* | Station Hill | Massive porphyritic granite |
| R27820* | 3 kms south of Bernborough Mine | Porphyritic rhyolite (Phase A) |
| | as above | as above |
| R27821 | as above | Intrusive ignimbrite (Phase B) |

University N.E. Geopeko Ltd.

| Rock Number | Thin Section Number | Polished Section Number | Locality | Lithology |
|-------------|---------------------|-------------------------|----------|-----------|
|-------------|---------------------|-------------------------|----------|-----------|

| | | | | |
|---------|--|--|---------------|--|
| R27822* | | | JUNO | |
| R27823* | | | 700 level | Basic dykes |
| R27824 | | | 700 level | Basic dyke |
| R27825* | | | 7/114/207 | Greywacke with shale inclusions |
| R27826* | | | 7/178/13-19.5 | Feldspathic greywacke |
| R27827 | | | 7/178/45-51 | Feldspathic Greywacke |
| R27828* | | | 8/159/59-65 | Coarse feldspathic greywacke |
| R27829* | | | 7/178/19.5-27 | Feldspathic greywacke-shale |
| R27830* | | | 7/178/27-31 | Chlorite-hematite-calcite bif. |
| R27831* | | | 7/178/31-36 | Chlorite-hematite-calcite bif. |
| R27832* | | | Surface DDH1 | Chlorite-hematite-calcite bif. |
| R27833 | | | 7/178/29 | Chlorite-hematite-calcite bif. |
| R27834 | | | 7/208/382 | Chlorite-hematite-calcite bif. |
| R27835 | | | 7/208/148 | Chlorite-hematite-calcite bif. |
| R27836 | | | 7/207/78 | Chlorite-hematite-calcite bif. |
| R27837 | | | 7/136/46 | Cherty siltstones |
| R27838* | | | 7/207/94 | Cherty siltstones |
| R27839 | | | Surface DDH1. | Crystal tuff ? |
| R27840 | | | 7/208/126 | Intraformational breccia |
| R27841 | | | 9/71/254 | Tuffaceous greywacke/shale |
| R27842 | | | 9/73/306 | Coarse tuffaceous greywacke |
| R27843 | | | 9/71/456 | Coarse tuffaceous greywacke |
| R27844 | | | 9/71/76 | Tuffaceous greywacke-shale |
| R27845 | | | 9/71/408 | Tuffaceous greywacke-shale |
| R27846 | | | 9/71/382 | Magnetite rich shale |
| R27847 | | | 9/71/110 | Tuffaceous greywacke shale |
| R27848 | | | 9/71/321 | Tuffaceous greywacke shale |
| R27849* | | | 7/25/44 | Altered bif. |
| R27850 | | | 700 level | Altered bif. |
| R27851 | | | 9/71/312 | Greywacke/shale |
| R27852* | | | 9/75/769 | Laminated magnetic shale |
| | | | 8/128/5 | Chloritised sediments adjacent to lode |

| University N.E. | | Geopeko Ltd. | | | |
|-----------------|------------------------|----------------------------|-----------|--|--|
| Rock Number | Thin Section Number | Polished Section Number | Locality | Lithology | |
| | | | | | |
| R27853* | | | 8/128/13 | Chloritised sediments adjacent to lode | |
| R27854*a | | | 8/128/21 | Chloritised sediments adjacent to lode | |
| R27855* | | | 8/128/29 | Chloritised sediments adjacent to lode | |
| R27856*a | | | 8/128/37 | Chloritised sediments adjacent to lode | |
| R27857* | | | 8/128/45 | Chloritised sediments adjacent to lode | |
| R27858*a | | | 8/128/53 | Chloritised sediments adjacent to lode | |
| R27859*a | | | 8/128/67 | Mineralised sediments | |
| R27860* | | | 8/128/65 | Mineralised sediments | |
| R27861*a | | GP 187 | 8/128/69 | Mineralised sediment | |
| R27862*a | | | 8/128/78 | Mineralised sediment | |
| R27863*a | | | 8/128/85 | Mineralised sediments | |
| R27864a | | | 8/128/80 | Mineralised sediments | |
| R27865*a | | | 8/128/93 | Magnetite-chlorite lode | |
| R27866*a | | | 8/128/101 | Magnetite-chlorite lode | |
| R27867*a | | | 8/128/109 | Magnetite-chlorite lode | |
| R27868 | | GP 189 | 8/128/113 | Magnetite-chlorite lode | |
| R27869*a | | | 8/128/117 | Magnetite-chlorite lode | |
| R27870*a | | | 8/128/126 | Magnetite-chlorite lode | |
| R27871*a | | | 8/128/133 | Magnetite-chlorite lode | |
| R27872*a | | | 8/128/141 | Magnetite-chlorite lode | |
| R27873*a | | | 8/128/149 | Talc-magnetite lode | |
| R27874*a | | | 8/128/157 | Talc-magnetite lode | |
| R27875*a | | | 8/128/165 | Dolomite lode | |
| R27876* | | GP 203 | 8/128/169 | Dolomite lode | |
| R27877*a | | | 8/128/173 | Dolomite lode | |
| R27878*a | | | 8/128/189 | Chloritised sediment | |
| R27879*a | | | 8/128/197 | Unaltered sediment | |
| R27880 | GT 703 | | 8/170/100 | Partially chloritised sediment | |
| R27881 | GT 863 | | 7/25/40 | Partially chloritised sediment | |
| R27882 | GT 613 | | 8/159/0-5 | Chloritised sediment | |
| R27883 | | GP 229 | 700 level | Magnetite-chlorite lode | |
| R27884a | | | 7/89/1 | Magnetite-chlorite lode | |

| University N.E. Rock Number | Geopeko Ltd. | | Locality | Lithology |
|-----------------------------------|------------------------|----------------------------|-----------|--|
| | Thin Section Number | Polished Section Number | | |
| R27885a | | | 7/89/31 | Magnetite-chlorite lode |
| R27886a | | | 7/89/59 | Magnetite-chlorite lode |
| R27887 | | GP 150 | 7/38/50 | Chalcopyrite ore |
| R27888 | | GP 168 | 7/38/98 | Mineralised sediment |
| R27889 | | GP 162 | 7/38/140 | Mineralised sediment |
| R27890 | | GP 171 | 7/38/155 | Mineralised sediment |
| R27891 | | GP 170 | 7/38/200 | Mineralised sediment |
| R27892 | | GP 151 | 7/176/172 | Mineralised sediment |
| R27893 | TS 880 | GP 167 | 7/36/108 | Magnetite-chlorite lode |
| R27894 | TS 872 | GP 156 | 7/36/117 | Talc-magnetite lode |
| R27895 | | GP 428 | 9/71/690 | Veins of mineralised sediment within hydrothermal channel |
| R27896 | | GP 407 | 9/73/845½ | Veins of mineralised sediment within hydrothermal channel |
| R27897 | | | 700 level | Talc-magnetite rock |
| R27898 | | | 700 level | Talc-magnetite rock |
| R27899* | | GP 137 | 7/167/21 | Talc-magnetite rock |
| R27900* | | GP 118 | 7/167/37 | Talc-magnetite rock |
| R27901* | | | 7/167/45 | Talc-magnetite rock |
| R27902* | | GP 130 | 7/167/69 | Talc-magnetite rock |
| R27903*a | | | 600 level | Banded dolomite |
| R27904* | | | 7/30/300 | Banded dolomite |
| R27905 | | | 600 level | Banded dolomite |
| R27906 | | | 700 level | Vein of banded dolomite |
| R27907a | | | 7/90/69 | Dolomite-jasper rock |
| R27908 | | | 7/89/5 | Magnetite-chlorite lode |
| GECKO MINE | | | | |
| R27909 | GT 1451 | | Anomaly 2 | Potassic greywacke/shale |
| | | | DDH22/484 | |
| R27910 | GT 1447 | | DDH22/317 | Intraformational breccia |
| R27911 | GT 1449 | | DDH22/411 | Intraformational breccia |
| R27912 | | | 2 level | Breccia-conglomerate |

University N.E.

Geopeko Ltd.

Rock
NumberThin Section
NumberPolished Section
Number

Locality

Lithology

| | | | | |
|--------|---------|--------|----------------|--|
| R27913 | GT 1374 | GP 501 | 2 level | Breccia-conglomerate |
| R27914 | GT 1380 | | 2 level | Breccia-conglomerate |
| R27915 | GT 1440 | | 2/9180/5/24 | Hematite shale |
| R27916 | | | Anomaly 1 | Hematite shale |
| R27917 | GT 1437 | | 2/9180/5/22/2 | Hematite shale |
| R27918 | GT 648 | | Anomaly 3 | Hematite shale |
| R27919 | GT 850 | | 2/9180/3/85 | Chloritised breccia-conglomerate |
| R27920 | TS 837 | GP 224 | 2/9180/3/171 | Altered breccia-conglomerate |
| R27921 | GT 1424 | GP 554 | 2/9180/8/6.8 | Altered breccia-conglomerate |
| R27922 | | | 2 level | Partially replaced breccia-conglomerate |
| R27923 | | | 2 level | Partially replaced breccia-conglomerate |
| R27924 | | | 2 level | Partially replaced breccia-conglomerate |
| R27925 | | | 2 level | Completely replaced breccia-conglomerate |
| R27926 | GT 1477 | | 4 level | Banded hematite-quartz lode |
| R27927 | | GP 249 | 2/9180/6/30 | Massive hematite with chalcopyrite |
| R27928 | | GP 549 | 2/9180/8/8.2 | Hematite-chlorite lode |
| R27929 | | GP 548 | 2/9180/8/17.5 | Massive hematite |
| R27930 | | GP | Anomaly 2 | Quartz-hematite |
| | | | DDH5/633 | |
| R27931 | | GP 212 | Anomaly 2 | Quartz-hematite |
| | | | DDH7/758 | |
| R27932 | | GP 256 | 2/9180/3/182.5 | Quartz-hematite |
| R27933 | | | 4 level | Massive magnetite-chalcopyrite |
| R27934 | | GP 247 | 2/9180/6/185 | Magnetite-chalcopyrite ore |
| R27935 | | GP 250 | 2/9180/6/141 | Chalcopyrite ore |
| R27936 | | GP 245 | 2/9180/6/95 | Magnetite-pyrite-chalcopyrite ore |
| R27937 | | GP 261 | 2/9180/6/162 | Magnetite-chalcopyrite ore |
| R27938 | | GP 248 | 2/9180/6/66 | Magnetite-chalcopyrite ore |
| R27939 | | GP 547 | 2/9180/8/30.8 | Magnetite-chalcopyrite ore |

| University N.E. Rock Number | Geopeko Ltd. | | Locality | Lithology |
|-----------------------------------|------------------------|----------------------------|-----------------|-------------------------------------|
| | Thin Section Number | Polished Section Number | | |
| R27940 | | GP 539 | 2/9180/8/39.3 | Magnetite-chalcoppyrite ore |
| R27941 | | GP 553 | 2/9180/8/5213 | Magnetite-bismuth ore |
| R27942 | | GP 541 | 2/9180/8/70.1 | Massive magnetite |
| R27943 | | GP ? | 2/9180/8/71.8 | Magnetite-bismuth ore |
| R27944 | | | Anomaly 2 | Magnetite-hematite lode |
| | | | DDH1/709 | |
| R27945 | | GP 220 | 2/9180/3/143 | Magnetite-chlorite-chalcoppyrite |
| R27946 | | GP 227 | 2/9180/3/139 | Magnetite-chalcoppyrite-pyrite |
| | | | GECKO ANOMALY 3 | |
| R27947* | | | H3/339 | Micro diorite |
| R27948* | | | H3/420 | Hornfelsed sediment |
| R27949* | | | H3/557 | Dolomite lode |
| R27950* | | | H3/610 | Dolomite lode |
| R27951* | | | H3/623 | Garnet-calcite rock |
| R27952*a | | | H3/749 | Magnetite-bornite ore |
| R27953 | | | H3/803 | Magnetite-calcite |
| R27954a | | | H3/835 | Magnetite-chalcoppyrite ore |
| R27956* | | GP 129 | H3/847 | Magnetite-calcite |
| R27957 | | | H3/766 | Magnetite-chalcoppyrite-bornite ore |
| R27958 | | | H3/905 | Magnetite-dolomite |
| R27959 | | | H3/942 | Chlorite-magnetite-dolomite |
| R27960 | | GP 143 | H3/964 | Magnetite-chalcoppyrite ore |
| R27961 | | | H3/1006 | Brecciated wall rock |
| R27962* | | | H3/929 | Chlorite-magnetite |
| | | | WARREGO ROCKS | |
| R27963* | | | 3 level | Quartzose slates & phyllites |
| R27964* | | | 3 level | Quartzose slates & phyllites |
| R27965* | | | 8/822/29/178 | Quartz-magnetite lode |
| R27966* | | | 8/822/29/207 | Magnetite-pyrite |
| R27967 | | | 8/822/29/243 | Massive magnetite |
| R27968* | | | 8/822/29/333 | Magnetite-chalcoppyrite |
| R27969* | | GP 280 | 8/822/29/342 | Bismuth lode with marcasite |

| University N.E. Rock Number | Geopeko Ltd. | | Locality | Lithology |
|-----------------------------------|------------------------|----------------------------|----------------|--|
| | Thin Section Number | Polished Section Number | | |
| R27970 | | GP 255 | 8/822/29/243 | Bismuth ore |
| R27971* | | | 8/822/29/382 | Quartz-magnetite |
| R27972* | | | 8/822/29/401 | Altered wall rock zone |
| R27973* | | | 8/822/29/415 | Tourmaline wall rock |
| R27974* | | | 8/822/29/280 | Magnetite-quartz |
| R27975 | | GP 296 | 6/824/10/196 | Magnetite-pyrite |
| R27976 | | GP 278 | 6/824/10/217 | Magnetite-pyrite |
| R27977 | | | 6 level | Magnetite-chalcoppyrite |
| R27978 | GT 1455 | GP ? | 9/816/22/52.8 | Magnetite-chlorite-quartz |
| R27979 | | GP 574 | 9/816/22/60.7 | Magnetite-chlorite |
| R27980 | GT 1468 | GP 573 | 9/816/22/65.1 | Magnetite-chlorite with gold |
| R27981 | | GP 587 | 9/816/22/89.8 | Magnetite-chlorite with chalcoppyrite |
| R27982 | | GP 583 | 9/816/22/92 | Magnetite-chlorite with chalcoppyrite |
| R27983* | | | 8/822/39/108 | Quartz-magnetite |
| R27984 | GT 1460 | | 8/816/22/115.2 | Muscovite alteration zone |
| R27985 | GT 1481 | | 8/816/22/116 | Andalusite alteration zone |
| R27986* | | | 8/822/29/171 | Chloritoid alteration zone |
| R31711* | | | 8/822/39/321 | Quartz porphyroid |
| R31712 | | | 9/816/22/118.5 | Altered quartz porphyroid |
| EXPLORER 46 | | | | |
| R31713 | GT 1357 | GP 491 | H4/3/307.7 | Typical chlorite-sericite shale |
| R31714 | GT 1360 | GP 494 | H4/3/353.7 | Hematite shale |
| R31715 | GT 1361 | GP 495 | H4/3/362.3 | Sandy calcite greywacke |
| R31716 | GT 1362 | GP 496 | H4/3/378.5 | Calcite greywacke |
| R31717 | GT 1364 | GP 498 | H4/3/383.3 | Sandy greywacke |
| R31718 | GT 1363 | | H4/3/383.2 | Coarse lithic wacke |
| R31719 | GT 1386 | GP 509 | H4/3/439 | Dolomite lode |
| R31720 | GT 1384 | GP 507 | H4/3/443 | Dolomite lode |
| R31721 | | GP 515 | H4/3/446 | Pyrite lode |
| R31722 | | GP 566 | H4/3/451.3 | Pyrite-magnetite lode |

| University N.E. Rock Number | Geopeko Ltd. | | Locality | Lithology |
|-----------------------------------|------------------------|----------------------------|--------------------------------------|----------------------|
| | Thin Section Number | Polished Section Number | | |
| R31723 | | GP 513 | H4/3/454.6 | Magnetite-chlorite |
| R31724 | | GP 511 | H4/3/457.6 | Bismuth ore |
| R31725 | GT 1385 | GP 508 | H4/3/471.6 | Calcite-pyrite |
| R31726 | | GP 411 | Juno, 9/71/14A | Tuffaceous greywacke |
| R31727 | GT 1227 | | Juno, 9/73/901 | Mineralised breccia |
| R31728 | GT 1225 | | Juno, 9/73/851 | Leached silt |
| R31729 | GT 1213 | GP 444 | Juno, 9/73/238 | Leached sediment |
| R31730 | GT 1289 | GP 457 | Juno, 9/69/516 | Mineralised breccia |
| R31731 | GT 1292 | GP 460 | Juno, 9/69/537 | Mineralised breccia |
| R31732 | | | 3/4 km north of Golden Forty Mine | Peko porphyry |
| R31733 | GT 1411 | | as above | Peko porphyry |
| R31734 | GT 1373 | | as above | Peko porphyry |
| R31735 | | | Juno, 7/207/87½ | Hematite shale |




Fig 1

This map will be reduced and simplified & include a locality map of Aust.

(interpretive).




Figure 1.1: - Regional Geology of the Tennant Creek Goldfield (interpretive).
 Compiled by Geopeko Ltd.